TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.

PRECAUTIONS

- 1. Handle with care, avoid splashing, contact with skin and eyes.
- 2. **DO NOT EAT, DRINK OR SMOKE WHEN USING THIS PRODUCT.**
- 3. Remove protective clothing on completion of spraying and wash hands and face thoroughly with water.
- 4. KEEP OUT OF REACH OF CHILDREN.
- 5. Store in original container and **KEEP UNDER LOCK AND KEY.**
- Dispose empty container, rinse, punctuate and bury in a safe place. **DO NOT** use for any other purpose.

SYMPTOMS OF POISONING

Mild irritation of eyes.

FIRST AID

Inhalation: In case of inhalation move the victim to fresh air. Seek medical advice if symptoms develop. **Eye contact**: Rinse immediately with plenty of water for 15-20 minutes: Remove contact lenses if available after the first 5 minutes, then continue rinsing. Seek medical advice if symptoms develop. **Skin contact**: Remove contaminated clothing. Wash immediately with plenty of water for 15-20 minutes. Seek medical advice if needed.

Ingestion: If swallowed, wash mouth with water provided the victim is conscious. Seek medical advice. **DO NOT** induce vomiting.

Call Drug and Toxicology Information Services on 0242307148 / 0786100098 for additional information on first aid.

NOTE TO PHYSICIAN

No specific antidote, treat symptomatically.

EMPTY CONTAINER DISPOSAL

Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Rinse the container 3 times with a volume of water equal to at least 25% of that of the container. Add the rinsate to the contents of the spray tank. Destroy the empty container by perforation and flattening. Place it in a secure disposal area and offer it for recycling. **DO NOT** use it for any other purpose.

DECONTAMINATION OF SPRAYER

Drain and rinse tank with CLEAN water, clean in the inside and outside. To clean the hoses and nozzles run CLEAN water through the system 3 times and spray out into a drain at the designated site for pesticide disposal.

WARRANTY


Although this insecticide has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label as well as by the method, time, and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the insecticide concerned due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier or registration holder in the event of any uncertainty.

COMPATABILITY

Don't add any surfactants with Wetcit® Duo. Buffers and acidifiers and most pesticides and foliar fertlizers are compatible with Wetcit® Duo. If in doubt regarding compatibility conduct compatibility tests in a smaller container and in the case of crop safety conduct tests on a small portion of the target crop before applying wholescale. Wetcit® Duo may interact with visible residues of previously applied products – therefore safety testing on a small portion of the crop is recommended. If in doubt, consult the technical representative of the Distributor.



INSECTICIDE



COMPOSITION: (mass/volume)

Chemical Group - UNE

WETCIT DUO is a soluble liquid insecticide for the control of red spider mite as indicated in specified crops.

Registered by Saiwit Holdings (Pvt) Ltd Stand 583 Pomona Industrial Park, Harare Telephone 08677005277

Manufactured by ORO AGRI, South Africa

DIRECTIONS FOR USE

USE ONLY AS DIRECTED.

EVEN DISTRIBUTION OF DROPLETS IS OF THE UTMOST IMPORTANCE FOR ALL APPLICATIONS. Ensure that both surfaces of the leaves (top and bottom) are thoroughly wet, up to just before the point of run-off. When used as a insecticide it is essential that the spray solution makes direct contact with the target organism during application.

MODE OF ACTION

INSECTICIDE: When applied at rates of 200ml / 100l of water and higher WETCIT DUO is a contact insecticide with no residual activity. Orange oil breaks down protective layers on mites. Physical disruption and breakdown of the protective layers occur only upon direct contact while the spray residue remains wet and results in the desiccation of the organism. ADJUVANT: At rates of 25-200ml / 100l of water WETCIT DUO drastically reduces the surface tension of water through the action of the non-ionic surfactant. This ensures even spreading of the spray mixture on the plant surface.

MIXING INSTRUCTIONS

Add WETCIT DUO at the end just before reaching the final volume in the spray tank, other products such as a buffer or acidifier must be added first, followed by the chemicals and then the WETCIT DUO.

DOSAGE INSTRUCTIONS

Crop	Pest	Application Rate	Remarks
Tomatoes	Red Spider Mite Tetranychus urticae	250 – 500 ml / 100 l of water 0.25 – 0.5 % solution 40 - 80 ml / 16 l of water 0.25 – 0.5 % solution	Start spraying at the first sign of infestation. Thorough wetting of both sides of the leaves must be always ensured. Do a small scale phytotoxicity test on varieties known to be sensitive or prone to phytotoxicity or on newly developed varieties. 3 Applications at 7-day intervals is recommended. DO NOT SPRAY TO THE POINT OF RUN-OFF. RECOMMENDED SPRAY VOLUME: Not less than 250 & / ha.

Re-entry: **DO NOT** enter treated areas within 6 hours of application.

Allow 6 hours between last application and harvest on tomatoes.

This product was tested under a protected environment.

IMPORTANT GUIDELINES

The surfactant properties of WETCIT DUO may result in dislodging residual spray material in the spray tank and hoses. Ensure that the spray tank and hoses are properly cleaned before applying WETCIT DUO.

DATE OF MANUFACTURE. BATCH NUMBER. EXPIRY DATE

