

PREV-AM[®]

READ THE ATTACHED PACKAGED LEAFLET BEFORE USE • KEEP OUT OF REACH OF CHILDREN AND ANIMALS

ACT / WET 36 / 1947 REG. No NAM N-AR 1303

PESTICIDE

A soluble liquid pesticide for the control of insects, mites, and diseases as indicated in specified crops.



WARNING

HAZARD STATEMENTS:

- May cause an allergic skin reaction
- Causes serious eye irritation
- Harmful if inhaled
- Harmful to aquatic life

PRECAUTIONARY STATEMENTS:

- Do not eat, drink or smoke when using this product
- Wash after use
- Avoid release to the environment
- Wear eye protection

ACTIVE INGREDIENTS

Orange oil

50 g / ℓ

REGISTRATION HOLDER:

Oro Agri SA (Pty) Ltd.

REG. No 2001/027414 / 07

POST 1 Henry Vos Close, ASLA Park, Strand, 7137

EMAIL info.ssa@rovensanext.com

WEB www.rovensanext.co.za

TEL +27 21 850 0667

EMERGENCY NUMBER: CHEMTREC SOUTH AFRICA 0-800 983 611

SEE CONTAINER FOR:

NET MASS / BATCH NO. / MANUFACTURING DATE

 **Rovensa
Next**



CAUTION • VERSIGTIG



WARNINGS

- MAY CAUSE AN ALLERGIC SKIN REACTION.
- CAUSES SERIOUS EYE IRRITATION.
- HARMFUL IF INHALED.
- HARMFUL TO AQUATIC LIFE.
- STORE IN A COOL, DRY PLACE AWAY FROM FOOD AND FEEDS.
- RE-ENTRY: Do not enter the treated area before the spray deposit has dried unless wearing protective clothing.
- Keep out of reach of children, uninformed persons, and animals.

PRECAUTIONS

- DO NOT EAT, DRINK OR SMOKE WHEN USING THIS PRODUCT.
- WASH AFTER USE.
- AVOID RELEASE TO THE ENVIRONMENT.
- WEAR EYE PROTECTION.
- IF SKIN IRRITATION OR RASH OCCURS: GET MEDICAL ADVICE/ ATTENTION.
- IF EYE IRRITATION PERSISTS: GET MEDICAL ADVICE/ ATTENTION.
- Wash contaminated clothing daily.
- Do not eat, drink, or smoke whilst applying or preparing the spray mix, or before washing your face and hands.
- Prevent drift of spray mist onto other crops, grazing, rivers, dams, and areas not under treatment.
- Clean applicator after use and dispose of water where it will not contaminate rivers, dams, grazing, and crops.
- Prevent contamination of food, feeds, drinking water, and eating utensils.
- Empty all pesticides from the container by placing it upside down over the spray tank and holding it there for at least 30 seconds.
- Triple rinse statement:
 - Fill empty container with one quarter of clean water
 - Shake for 30 seconds
 - Decant into spray tank for 30 seconds
- NEVER USE THE CONTAINER FOR ANY OTHER PURPOSE. Puncture the rinsed container to render it useless and send it to a CropLife SA-approved recycler. PET containers cannot be recycled and must be incinerated after cleaning.

RELEVANT SUBSTANCES

- Alcohols, C10-16, ethoxylated, sulphates, sodium salts > 10 %; Alcohol ethoxylate (non-ionic surfactant) > 5 %; Orange oil > 5 %; Sodium hydroxide > 0.5 %

DISCLAIMER: Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label as well as by the method, time and accuracy of the application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration.

Consult the supplier or registration holder in the event of any uncertainty.

CONSULT THE SUPPLIER OR REGISTRATION HOLDER IN THE EVENT OF ANY UNCERTAINTY.

FIRST AID

• IN CASE OF INHALATION:

If breathed in, move person into the fresh air. If not breathing, give artificial respiration. In case of discomfort seek medical attention.

• IN CASE OF SKIN CONTACT:

Remove contaminated clothing; wash affected area with water for 15 minutes. In case of discomfort or irritation seek medical attention.

• IN CASE OF EYE CONTACT

Flush eyes thoroughly with water for several minutes. Remove contact lenses after the initial 1-2 minutes and continue flushing for several additional minutes. Seek medical advice if symptoms or irritation persist.

• IN CASE OF INGESTION:

Do NOT induce vomiting. Never give anything by mouth to an unconscious person. Rinse mouth with water. Seek medical attention if you feel unwell.

MODE OF ACTION

- **PREV-AM®** is a contact insecticide/fungicide with no residual activity and as such should be used as a direct contact application at first sight of insects or disease.

COMPATABILITY

- Experience has shown that **PREV-AM®** is compatible with most crop protection products and foliar fertilizers.
- Do not mix **PREV-AM®** with other products unless prior tests (including small-scale phytotoxicity tests) have shown the combination to be physically compatible, effective, and non-injurious to the target crop under local conditions.
- A phytotoxicity test must also be done if visible residues of previously applied crop protection chemicals or foliar fertilizers are present on fruit, as these may be loosened by the **PREV-AM®** application and washed down to collect in a droplet, thus increasing the risk of phytotoxic blemishes on the fruit.
- If in doubt, consult the technical representative of the distributor.
- Ensure that the spray tank and hoses are properly cleaned inside before applying **PREV-AM®**, as the product may loosen residues and adsorbed materials of old products which could contaminate the spray mixture.
- In the event of any of these prescriptions not having been followed, the Manufacturer and its Distributors accept no responsibility for any possible damage or harmful consequences that may ensue.
- Although **PREV-AM®** has been exempted from maximum residue levels (MRL) and has no pre-harvest withholding period, consult the labels of tank-mix products to ensure adherence to withholding periods between the last application and harvest periods of these products.

PRODUCT COMPOSITION AND METHOD PATENTED · PRODUKSTAMESTELLING EN -METODE GEPATENTEER ZA200405010 / ZA200703565

Oro Agri SA (Pty) Ltd. (THE COMPANY) WARRANTS THAT THE PRODUCT CONFORMS TO SPECIFICATIONS ON THE LABEL AND IS REASONABLY FIT FOR THE INTENDED PURPOSE REFERRED TO ON THE LABEL. THE COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. THE COMPANY RECOMMENDS THAT THE USER TESTS THIS PRODUCT ON A PORTION OF THE CROP TO DETERMINE SUITABILITY FOR THE INTENDED USE.

GROUND APPLICATION

ONLY SUITABLE APPLICATION EQUIPMENT MUST BE USED

- FOR THE APPLICATION OF SPRAYS IN TUNNEL GROWING OR OPEN FIELD CROPS, THE EVEN DISTRIBUTION OF THE DROPLETS IS OF THE UTMOST IMPORTANCE.
- Care must be taken to ensure that both surfaces of the leaves are thoroughly wet, up to just before the point of run-off. DO NOT SPRAY UNTIL DRIPPING WET, AS THIS MIGHT CAUSE THE COLLECTION OF TOO MUCH PRODUCT IN PLACES.
- PREV-AM® only acts as a contact insecticide/fungicide – it is, therefore, essential that the spray solution make direct contact with the insect (in any stage of its development cycle) or with the disease organism during application.
- Applications by suitable tractor-mounted sprayers must ensure that a thorough wetting of both the top and bottom surfaces of the leaves is achieved.
- The use of conventional tractor boom-sprayers usually does not achieve adequate wetting of the bottoms of the leaves. The use of a mist-blower, tractor sprayer, or any other sprayer with which the spray mist could be blown into the crop or leaves (where an oscillating mechanism will be an added advantage), is recommended.

DOSAGE RATES

CUCURBITS

PEST / DISEASE	APPLICATION RATE	REMARKS
Whitefly <i>Bemisia tabaci</i>	0.5 l PREV-AM® per 100 l water 0.5% solution	<ul style="list-style-type: none"> • Applications should start within three weeks after emergence of seedlings and continued as a weekly spray programme (7-day intervals) as long as needed. • Thorough wetting of both sides of the leaves must be ensured at all times.
Powdery Mildew <i>Sphaerotheca fuliginea</i>	0.5 l PREV-AM® per 100 l water 0.5% solution	<ul style="list-style-type: none"> • DO NOT SPRAY TO THE POINT OF RUN-OFF. • RECOMMENDED SPRAY VOLUME: 500 l/ha

CITRUS

PEST / DISEASE	APPLICATION RATE	REMARKS
<p>Citrus Black Spot</p>	<p>0.1 l PREV-AM[®] + 150 g <i>mancozeb 800WP</i> + 20-30 ml <i>azoxystrobin 250SC</i> OR 10 ml <i>pyraclostrobin 250EC</i> per 100 l water</p>	<ul style="list-style-type: none"> A preventative programme for the control of citrus black spot should commence with a <i>mancozeb 800WP</i> application (as registered) at fruit set. Apply the first PREV-AM[®] + <i>mancozeb 800WP</i> + <i>azoxystrobin 250SC</i> or <i>pyraclostrobin 250EC</i> application no later than 24 days after the last <i>mancozeb 800WP</i> application, and follow up with a second PREV-AM[®] + <i>mancozeb 800WP</i> + <i>azoxystrobin 250SC</i> or <i>pyraclostrobin 250EC</i> application 6 weeks later. Complete the program with a <i>mancozeb 800WP</i> application (as registered) no later than 6 weeks after the second PREV-AM[®] + <i>mancozeb 800WP</i> + <i>azoxystrobin 250SC</i> or <i>pyraclostrobin 250EC</i> application. Apply as high volume applications to ensure proper coverage of fruit to be protected, and use the higher rate of <i>azoxystrobin 250SC</i> if conditions are favourable for black spot development or high infection levels were experienced the previous season. NOTE: Consult the <i>mancozeb 800WP</i>, <i>azoxystrobin 250SC</i> or <i>pyraclostrobin 250EC</i> labels for details.
<p><i>Guignardia citricarpa</i></p>	<p>0.1 l PREV-AM[®] + 150 g <i>mancozeb 800WP</i> per 100 l water</p>	<ul style="list-style-type: none"> Apply not later than 24 days after the <i>mancozeb 800WP</i> application at petal drop (as registered) and repeat 4-5 times at 24 day intervals. NOTE: Consult the <i>mancozeb 800WP</i> label for details.
	<p>0.1 l PREV-AM[®] + 200 g <i>mancozeb 800WP</i> + 50 g <i>benomyl 500WP</i> per 100 l water</p>	<ul style="list-style-type: none"> Only use <i>benomyl 500WP</i> in orchards where benzimidazole resistance is absent. Precede the PREV-AM[®] + <i>mancozeb 800WP</i> + <i>benomyl 500WP</i> application with an application of <i>mancozeb 800WP</i> at 200 g / 100 liters of water (as registered) in the last week of October. Apply one PREV-AM[®] + <i>mancozeb 800WP</i> + <i>benomyl 500WP</i> application as a full cover spray between the 1st and 20th of December. If obtaining sufficient fruit cover is a concern, two sprays of PREV-AM[®] (0.1 l) + <i>mancozeb 800WP</i> (200g) + <i>benomyl 500WP</i> (25g / 100l water) can be applied during the last half of November and the first week of January respectively. NOTE: Consult the <i>mancozeb 800WP</i> and <i>benomyl 500WP</i> labels for details.

PEST / DISEASE	APPLICATION RATE	REMARKS
<p>Citrus Thrips <i>Scirtothrips aurantii</i></p>	<p>0.2ℓ PREV-AM[®] per 100ℓ water 0.2% solution</p>	<ul style="list-style-type: none"> Apply between 80% and 100% petal drop as preventative application Since PREV-AM[®] is not restricted by a pre-harvest interval or maximum residue level, the product can be used as a late corrective application up to harvest, once fruit has fully coloured up.
<p>Red Scale <i>Aonidiella aurantii</i></p>	<p>0.2 - 0.4ℓ PREV-AM[®] per 100ℓ water 0.2 - 0.4% solution</p>	<ul style="list-style-type: none"> The use of PREV-AM[®] for the control of Red Scale should only commence once fruit has fully coloured up.
<p>Citrus Red Mite <i>Panonychus citri</i></p>	<p>0.2ℓ PREV-AM[®] per 100ℓ water 0.2% solution</p>	<ul style="list-style-type: none"> Apply when the first signs of a red mite infestation is observed. A repeat application 5 to 7 days after the first is recommended.

Do not apply PREV-AM[®] on Citrus crops at rates exceeding 100 mℓ/hℓ in the period from 100% petal fall up to and including full colour development.

PEST / DISEASE	APPLICATION RATE	REMARKS
Mealybug <i>Planococcus ficus</i>	0.5 ℓ PREV-AM [®] per 100 ℓ water 0.5% solution	<ul style="list-style-type: none"> • Monitor for mealybug movement and start spray application against mealybug as soon as crawlers are observed on shoots or leaves. Repeat application after 7 days if further crawler movement is observed. • Ants must be controlled separately before mealybug control is done. • RECOMMENDED SPRAY VOLUME: 750 ℓ/ha to 1000 ℓ/ha, depending on vine size.
Powdery Mildew <i>Uncinula necator</i>	0.25 - 0.5 ℓ PREV-AM [®] per 100 ℓ water 0.25 - 0.5% solution	<ul style="list-style-type: none"> • A programme for the control of powdery mildew should commence when the longest shoots reach a length of 2-5 cm. Use conventional residual fungicides up to flowering. • The optimal window for use of PREV-AM[®] will be after flowering up to harvest or during the period when mealybug crawler movement is expected (see mealybug recommendation). • Apply PREV-AM[®] at 7-10 day intervals as part of a powdery mildew control program, alternating with products with different modes of action. PREV-AM[®] has a non-specific contact action and is therefore suitable for use in an anti-resistance programme against powdery mildew. • Preferably use a disease forecasting model as well as disease history of the target vineyard as a guideline to assess risk of disease. If this information indicates low disease pressure and low risk, or no symptoms are visible, use the 0.25% dosage rate. When higher disease pressure and higher risk is indicated, or symptoms are already visible, use the 0.5% dosage rate. • Should conditions be favourable for the infection of clusters with green berries, PREV-AM[®] should be alternated or combined with a suitable systemic fungicide to assist in the protection of the clusters. • RECOMMENDED SPRAY VOLUME: 500 ℓ/ha to 1200 ℓ/ha, depending on vine size.
Downy Mildew <i>Plasmopara viticola</i>	0.5 ℓ PREV-AM [®] per 100 ℓ water 0.5% solution	<ul style="list-style-type: none"> • A programme for the control of downy mildew should commence when the longest shoots reach a length of 10 cm. Use conventional residual fungicides for preventative control of downy mildew. • If PREV-AM[®] is applied at 500 mℓ/100ℓ for mealybug or powdery mildew, this spray will also control downy mildew which may be present at the time. • If infection periods for downy mildew are expected after application of PREV-AM[®], it should be combined with a preventative residual fungicide as registered. • Ensure thorough wetting of underside of leaves. • RECOMMENDED SPRAY VOLUME: 750 ℓ/ha to 1000 ℓ/ha, depending on vine size.

LETTUCE**WESTERN AND SOUTH CAPE REGIONS ONLY**

PEST / DISEASE	APPLICATION RATE	REMARKS
Downy Mildew <i>Plasmopara viticola</i>	0.5 l PREV-AM [®] per 100 l water 0.5% solution	<ul style="list-style-type: none"> • Start applying at first signs of disease. • A spray programme with 3 - 7 day intervals is recommended. • Thorough wetting of both sides of the leaves must be ensured at all times. • DO NOT SPRAY TO THE POINT OF RUN-OFF. • In event of extreme diseases infections levels, the addition of a suitable residual fungicide (as registered) will be required. • RECOMMENDED SPRAY VOLUME: Not less than 500 l/ha

ORNAMENTALS AND ROSES

PEST / DISEASE	APPLICATION RATE	REMARKS
Thrips spp. <i>Thripidae spp.</i>		
Aphids <i>Macrosiphum rosae</i>		<ul style="list-style-type: none"> • A spray programme with 3 - 7 day intervals is recommended.
Red Spider Mite <i>Tetranychus urticae</i>	0.5 l PREV-AM [®] per 100 l water 0.5% solution	<ul style="list-style-type: none"> • Thorough wetting of both sides of the leaves must be ensured at all times. • Do a small scale phytotoxicity test on varieties known to be sensitive or prone to phytotoxicity or on newly developed varieties.
Whitefly <i>Bemisia tabaci</i>		<ul style="list-style-type: none"> • DO NOT SPRAY TO THE POINT OF RUN-OFF.
Leaf Hoppers <i>Jassid spp.</i>		<ul style="list-style-type: none"> • RECOMMENDED SPRAY VOLUME: Not less than 250 l/ha
Powdery Mildew <i>Sphaerotheca fuliginea</i>		

TOMATOES

PEST / DISEASE	APPLICATION RATE	REMARKS
Whitefly <i>Bemisia tabaci</i>	0.5 l PREV-AM® per 100 l water 0.5% solution	<ul style="list-style-type: none"> Application should start within three weeks after transplanting and continued as a weekly spray programme (7-day intervals) as long as needed. Ensure thorough wetting of both sides of the leaves at all times. DO NOT SPRAY TO THE POINT OF RUN-OFF. This treatment for White fly will enhance the control of early and/or late blight, when combined in a tank mixture with a registered fungicide for blight control. The treatment for whitefly will also control aphids and red spider mites present during an application. RECOMMENDED SPRAY VOLUME: 500 l/ha, from flowering onwards.

RATES FOR BACKPACK SPRAYERS

PREV-AM®	BACKPACK VOLUME		
Dose per 100 liters water	10 liter	16 liter	20 liter
100 ml	10 ml	16 ml	20 ml
200 ml	20 ml	32 ml	40 ml
250 ml	25 ml	40 ml	50 ml
500 ml	50 ml	80 ml	100 ml

PREV-AM®

LEES DIE AANGEHEGTE ETIKET VOOR GEBRUIK • HOU BUIITE BEREIK VAN KINDERS EN DIERE

ACT / WET 36 / 1947 REG. No NAM N-AR 1303

PLAAGDODER

'n Oplosbare vloeistof plaagdoder vir die beheer van insekte, myte en siektes soos aangedui in gespesifiseerde gewasse.



WAARSKUWING

GEVAARSTELLINGS:

- Mag 'n allergiese velreaksie veroorsaak
- Veroorsaak ernstige oogirritasie
- Skadelik indien ingesam
- Skadelik vir waterlewe

VOORSORGSTELLINGS:

- Moenie eet, drink of rook wanneer hierdie produk gebruik word nie
- Was na gebruik
- Vermo vrylating na die omgewing
- Dra oogbeskerming

AKTIEWE BESTANDELE

Lemoenolie

50 g / ℓ

REGISTRASIE HOUER:

Oro Agri SA (Pty) Ltd.

REG. No 2001/027414 / 07

POS 1 Henry Vos Close, ASLA Park, Strand, 7137

E-POS info.ssa@rovensanext.com

WEB www.rovensanext.co.za

TEL +27 21 850 0667

EMERGENCY NUMBER: CHEMTRAC SOUTH AFRICA 0-800 983 611

SIEN HOUER VIR:

NETTO MASSA / LOT NO. / VERVAARDIGINGSDATUM

 **Rovensa
Next**



CAUTION • VERSIGTIG



WAARSKUWINGS

- MAG 'N ALLERGIESE VELREAKSIE VEROORSAAK.
- VEROORSAAK ERNSTIGE OOGIRRITASIE.
- SKADELIK INDIEN INGESAEM.
- SKADELIK VIR DIE WATERLEWE.
- BERG OP 'N KOEL, DROË PLEK WEG VAN KOS EN VOERE.
- HERBETREDING: Moenie die behandelde area betree voordat die spuitneerslag droog is nie, tensy beskermende klere gedra word.
- Hou buite bereik van kinders, ongeligte persone en diere.

VOORSORGMATREËLS

- MOENIE EET, DRINK OF ROOK WANNEER HIERDIE PRODUK GEBRUIK WORD NIE.
- WAS NA GEBRUIK.
- VERMY VRYSTELLING NA DIE OMGEWING.
- DRA OOGBESKERMING.
- INDIEN VEL IRRITASIE OF UITSLAG VOORKOM: KRY MEDIESE RAAD/ AANDAG.
- INDIEN OOG-IRRITASIE AANHOU: KRY MEDIESE RAAD/ AANDAG.
- Was besoedelde klere daaglik.
- Moenie eet, drink of rook terwyl die spuitmengsel toegedien of voorberei word nie, of voordat u gesig en hande gewas is nie.
- Voorkom wegdrywing van spuitnewel na ander gewasse, weiding, riviere, damme en gebiede wat nie behandel word nie.
- Maak spuitapparaat skoon na gebruik en doen weg met water waar dit nie riviere, damme, weiding en gewasse sal besoedel nie.
- Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.
- Maak alle plaagdoders uit die houer leeg deur dit onderstebo oor die spuitenk te plaas en dit vir minstens 30 sekondes daar te hou.
- Drievoudige spoelverklaring:
 - Vul lee houer 'n kwart met skoon water
 - Skud vir 30 sekondes
 - Laat in spuitenk uitloop vir 30 sekondes
- MOET NOOIT DIE HOUER VIR ENIGE ANDER DOEL GEBRUIK NIE. Kap die uitgespoelde houer vol gate om dit nutteloos te maak en stuur dit na 'n CropLife SA-geodekeurde herwinner. PET-houers kan nie herwin word nie en moet na skoonmaak verbrand word.

ALHOEWEL HIERDIE MIDDEL OMVATTEND ONDER 'N GROOT VERSKEIDENHEID TOESTANDE GETOETS IS, WAARBORG DIE REGISTRASIEHOUER NIE DAT DIT ONDER ALLE TOESTANDE DOELTREFFEND SAL WEES NIE. AANGESIEN DIE WERKING EN EFFEK DAARVAN BEÏNVOED KAN WORD DEUR FAKTORE SOOS ABNORMALE KLIMAATS- EN BERGINGSTOESTANDE, KWALITEIT VAN VERDUNNINGSWATER, VERENIGBAARHEID MET ANDER STOWWE WAT NIE OP DIE ETIKET AANGEDUI IS NIE. SOWEL AS DIE METODE. TYD EN AKKURATEID VAN TOEDIENING. VERDER AANVAAR DIE REGISTRASIEHOUER NIE VERANTWOORDELIKHEID VIR SKADE AAN GEWASSE, PLANTEGROEI, DIE OMGEWING OF VIR NADIELIGE EFFEK OP MENS EN DIER OF VIR 'N GEBREK AAN PRESTASIE VAN DIE BETROKKE MIDDEL AS GEVOLG VAN DIE VERSIJM DEUR DIE GEBRUIKER OM ETIKETAANWYSINGS NA TE KOM OF AS GEVOLG VAN DIE ONTSTAAN VAN TOESTANDE WAT NIE KRAGTENS DIE REGISTRASIE VOORSIEN KON WORD NIE.

RAADPLEEG DIE VERSKAFFER OF REGISTRASIEHOUER IN GEVAL VAN ENIGE ONSEKERHEID.

RELEVANTE STOWWE

- Alcohols, C10-16, ethoxylated, sulphates, sodium salts > 10 %; Alcohol ethoxylate (non-ionic surfactant) > 5 %; Orange oil > 5 %; Sodium hydroxide > 0.5 %

NOODHULP

• IN DIE GEVAL VAN INASEMING:

Indien ingeasem, beweeg persoon in die vars lug. Indien nie asemhaal nie, gee kunsmatige asemhaling. In geval van ongemak soek mediese hulp.

• IN GEVAL VAN VELKONTAK:

Verwyder besmette klere; was geaffekteerde area met water vir 15 minute. In geval van ongemak of irritasie soek mediese hulp.

• IN GEVAL VAN OOGKONTAK:

Spoel oë deeglik uit met water vir 'n paar minute. Verwyder kontaklense na die aanvanklike 1-2 minute en gaan voort om vir 'n paar ekstra minute te spoel. Soek mediese advies indien simptome of irritasie voortduur.

• IN GEVAL VAN INNAME:

MOENIE braking veroorsaak nie. Moet nooit iets per mond aan 'n bewustelose persoon gee nie. Spoel mond uit met water. Soek mediese hulp as jy ongesteld voel.

METODE VAN WERKING

- **PREV-AM®** is 'n kontak insekdoder/swamdoder met geen oorbywende aktiwiteit nie en moet as sodanig gebruik word as 'n direkte kontak toediening met die eerste oogopslag van insekte of siektes.

VERENIGBAARHEID

- Ervaring het getoon dat **PREV-AM®** met die meeste gewasbeskermingsprodukte en blaarbemesting versoenbaar is.
- Moenie **PREV-AM®** met ander produkte meng nie, tensy vorige toetse (insluitend kleinskaalse fitotoksiteitstoetse) getoon het dat die kombinasie fisies versoenbaar, doeltreffend en nie-skadelik vir die teikengewas onder plaaslike toestande is nie.
- 'n Fitotoksiteitstoets moet ook gedoen word indien sigbare oorbyfsels van voorheen toegediende gewasbeskermingsmiddels of blaarbemesting op vrugte aanwesig is, aangesien dit deur die **PREV-AM®** toediening losgemaak kan word en afgespoel kan word om in 'n druppel te versamel, en sodoende die risiko van fitotoksies verhoog vlekke op die vrugte.
- Indien jy twyfel, raadpleeg die tegniese verteenwoordiger van die verspreider.
- Maak seker dat die spuittenk en slange behoorlik binne skoongemaak is voordat **PREV-AM®** toegedien word, aangesien die produk residu en geabsorbeerde materiale van ou produkte wat die spuitmengsel kan besoedel, kan losmaak.
- In die geval waar enige van hierdie voorskrifte nie gevolg is nie, aanvaar die Vervaardiger en sy Verspreiders geen verantwoordelikheid vir enige moontlike skade of skadelike gevolge wat mag intree nie.
- Alhoewel **PREV-AM®** vrygestel is van maksimum residuvlakke (MRL) en geen voor-oes-onthoudingsperiode het nie, raadpleeg die etikette van tenkmengselprodukte om te verseker dat die onthoudingsperiodes tussen die laaste toediening en oesperiodes van hierdie produkte nagekom word.

GRONDTOEDIENING

SLEGS GEPASTE TOEDIENINGSTOERUSTING MOET GEbruik WORD

- **VIR DIE TOEDIENING VAN BEDIENINGE IN TONNELGROEIENDE OF OOPLAND GEWASSE IS DIE ELIGE VERSPREIDING VAN DIE DRUPPELS VAN DIE GROOTSTE BELANG.**
- Sorg moet gedra word om te verseker dat beide oppervlaktes van die blare deeglik nat is, tot net voor die punt van afloop.
- **MOENIE SPUIT TOT DRUPPEND NAT NIE, AANGESIEN DIT DIE VERSAMELING VAN TE VEEL PRODUK OP PLEKKE KAN VEROORSAAK.**
- **PREV-AM®** dien slegs as 'n kontak insekdoder/swamdoder – dit is dus noodsaaklik dat die spuitoplossing direkte kontak maak met die insek (in enige stadium van sy ontwikkelingsiklus) of met die siekte-organisme tydens toediening.
- Toedienings deur geskikte trekkergemonteerde spuite moet verseker dat 'n deeglike benatting van beide die boonste en onderste oppervlaktes van die blare verkry word.
- Die gebruik van konvensionele trekkerbalkspuite bewerkstellig gewoonlik nie voldoende benatting van die onderkant van die blare nie. Die gebruik van 'n misblaser, trekkerspuit, of enige ander spuit waarmee die spuitnewel in die gewas of blare ingeblaas kan word (waar 'n ossillerende meganisme 'n bykomende voordeel sal wees), word aanbeveel.

DOSSISE

BLAARSLAAI

SLEGS WES- EN SUID-KAAP GEBIEDE

PLAAG / SIEKTE	TOEDIENINGSHOEVEELHEID	OPMERKINGS
<p>Donsskimmel <i>Plasmopara viticola</i></p>	<p>500 ml PREV-AM® per 100 l water 0.5% spuitoplossing</p>	<ul style="list-style-type: none"> • Begin spuit sodra eerste tekens van siekte verskyn. • 'n Spuitprogramme met 3 - 7 dae tussenposes word aanbeveel. • Deeglike benatting van beide die bo- en onderkant van blare moet te alle tye verseker word. • Onder uitsonderlike siektedruk-toestande, sal die byvoeging van 'n geregistreerde swamdoder met residuele werking benodig word. • BESPUIT TOT NET VOORDAT AFLOOP PLAASVIND. • Let noukeurig op die voorskrifte van die VERENIGBAARHEID paragrawe. • AANBEVOLE SPUITVOLUME: Nie minder as 500 l/ha nie.

RANKGEWASSE

PLAAG / SIEKTE	TOEDIENINGSHOEVEELHEID	OPMERKINGS
Witvlieg <i>Bemisia tabaci</i>	500 ml PREV-AM [®] per 100 l water 0.5% spuitoplossing	<ul style="list-style-type: none">• Toediening moet begin binne drie weke nadat saailinge begin opkom en voortgesit word met 'n weeklikse spuitprogramme (7 dae tussenposes), vir so lank as nodig.• Deeglike benatting van beide die bo- en onderkant van blare moet te alle tye gedoen word.• MOENIE BESPUIT TOT PUNT VAN AFLOOP NIE.• Let noukeurig op die voorskrifte van die VERENIGBAARHEID paragrawe.• AANBEVOLE SPUITVOLUME: 500 l/ha
Poeieragtige Skimmel <i>Sphaerotheca fuliginea</i>		

SIERPLANTE EN ROSE

PLAAG / SIEKTE	TOEDIENINGSHOEVEELHEID	OPMERKINGS
Blaaspootjies <i>Thripidae spp.</i>	500 ml PREV-AM [®] per 100 l water 0.5% spuitoplossing	<ul style="list-style-type: none">• 'n Spuitprogram met 3 - 7 dae intervalle word aanbeveel.• Deeglike benatting van beide die bo- en onderkant van blare te alle tye verseker.• BESPUIT TOT NET VOORDAT AFLOOP PLAASVIND.• Let noukeurig op die voorskrifte van die VERENIGBAARHEID paragrawe.• AANBEVOLE SPUITVOLUME: Nie minder as 250 l/ha nie.
Plantluise <i>Macrosiphum rosae</i>		
Roospinmyte <i>Tetranychus urticae</i>		
Witvlieg <i>Bemisia tabaci</i>		
Bladspringers <i>Jassid spp.</i>		
Poeieragtige Skimmel <i>Sphaerotheca fuliginea</i>		

SITRUS**PLAAG / SIEKTE****TOEDIENINGSHOEVEELHEID****OPMERKINGS**

Blaaspotjies
Scirtothrips aurantii

200 mℓ **PREV-AM**[®]
per 100 ℓ water
0.2% spuitplossing

- Dien tydens 80% en 100% blomblaarval toe as voorkomende behandeling.
- Aangesien die gebruik van **PREV-AM**[®] nie deur 'n voor-oes onthoudingsperiode of maksimum residue vlakke beperk word nie, kan die produk as laat korrektiewe bespuiting op vrugte na volle kleur ontwikkeling tot oestyd gebruik word.

Rooidopluis
Aonidiella aurantii

200 - 400 mℓ **PREV-AM**[®]
per 100 ℓ water
0.2 - 0.4% spuitplossing

- Beperk die gebruik van **PREV-AM**[®] vir die beheer van rooi dopluis tot die periode na volle kleur ontwikkeling van die vrugte tot en met oes.

Sitrus
Rooi Myte
Panonychus citri

200 mℓ **PREV-AM**[®]
per 100 ℓ water
0.2% spuitplossing

- Dien toe wanneer die eerste tekens van 'n myt infestasië opgemerk word.
- 'n Opvolgbespuiting 5 tot 7 dae na die eerste word aanbeveel.
- Let noukeurig op die voorskrifte van die VERENIGBAARHEID paragrafe.

PLAAG / SIEKTE

TOEDIENINGSHOEVEELHEID

OPMERKINGS

Sitrus Swartvlek

Guignardia citricarpa

100 ml **PREV-AM**® +
150g *mankoseb 800WP* +
20-30ml *asoksiestrobien 250SC*
OF
10ml *piraklostrobien 250EC*
per 100 l water

- 'n Voorkomende program vir die beheer van sitrus swartvlek behoort 'n aanvang te neem met 'n *mankoseb 800WP* bespuiting (soos geregistreer) by vrugset.
- Dien die eerste **PREV-AM**® + *mankoseb 800WP* + *asoksiestrobien 250SC* of *piraklostrobien 250EC* toediening nie later as 24 dae na die laaste *mankoseb 800WP* toediening toe.
- Volg op met 'n tweede **PREV-AM**® + *mankoseb 800WP* + *asoksiestrobien 250SC* of *piraklostrobien 250EC* toediening 6 weke later.
- Voltooi die program met 'n *mankoseb 800WP* toediening (soos geregistreer) nie later as 6 weke na die tweede **PREV-AM**® + *mankoseb 800WP* + *asoksiestrobien 250SC* of *piraklostrobien 250EC* toediening.
- Dien as hoë volume toedienings toe ten einde volledige dekking van vrugte wat beskerm moet word, te verseker en gebruik die hoër *asoksiestrobien 250SC* dosis as toestande gunstig is vir die ontwikkeling van swartvlek, of indien hoë siektedruk die vorige seisoen ondervind is.
- Let noukeurig op die voorskrifte van die VERENIGBAARHEID paragrawe.
- NOTA: Raadpleeg die *mankoseb 800WP* en *asoksiestrobien 250SC* of *piraklostrobien 250EC* etikette vir detail.

100 ml **PREV-AM**® +
150g *mankoseb 800WP*
per 100 l water

- Dien toe nie later as 24 dae na die aanvanklike *mankoseb 800WP* toediening by blaarval (soos geregistreer) en herhaal 4-5 keer met 24 dae intervalle.
- NOTA: Raadpleeg die *mankoseb 800WP* etiket vir detail.

100 ml **PREV-AM**® +
200g *mankoseb 800WP* +
50g *benomil 500WP*
per 100 l water

- Gebruik *benomil 500WP* slegs in boorde waar bensimidiasool weerstand afwesig is.
- Gaan die **PREV-AM**® + *mankoseb 800WP* + *benomil 500WP* bespuiting vooraf met 'n *mankoseb 800WP* toediening teen 200g per 100 l water (soos geregistreer) in die laaste week van Oktober.
- Dien een **PREV-AM**® + *mankoseb 800WP* + *benomil 500WP* toediening as voldekbepuiting toe tussen die 1ste en 20ste Desember.
- Indien kommer bestaan dat onvoldoende dekking met bespuitings verkry word, kan twee **PREV-AM**® (0.1 l) + *mankoseb 800WP* (200g) + *benomil 500WP* (25g per 100 l water) bespuitings gedurende die laaste helfte van November en weer die eerste week Januarie gedoen word.
- NOTA: Raadpleeg die *mankoseb 800WP* en *benomil 500WP* etikette vir detail.

Moenie 'n **PREV-AM**® op sitrus gewasse toedien teen dosisse wat 100 ml/hl oorskry vir die periode van 100% blomblaarval tot en met volle kleur ontwikkeling nie.

TAMATIES

PLAAG / SIEKTE	TOEDIENINGSHOEVEELHEID	OPMERKINGS
Witvlieg <i>Bemisia tabaci</i>	500 mℓ PREV-AM® per 100 ℓ water 0.5% spuitoplossing	<ul style="list-style-type: none"> • Toediening moet binne drie weke ná oorplanting begin en voortgesit word met 'n weeklikse spuitprogramme (7-dae tussenposes), vir so lank as nodig. • Deeglike benatting van beide die bo- en onderkant van blare te alle tye verseker word. • MOENIE BESPUIT TOT PUNT VAN AFLOOP NIE. • Hierdie behandeling vir witvlieg sal die beheer van vroeë- of laatros verbeter, indien dit gekombineer word in tenkmengsel met geregistreerde swamdoders teen hierdie siektes. • Die behandeling vir witvlieg sal ook plantluus of roospinmyt beheer indien teenwoordig tydens bespuiting. • Let noukeurig op die voorskrifte van die VERENIGBAARHEID paragrawe. • AANBEVOLE SPUITVOLUME: 500 ℓ/ha word vanaf die blomstadium en verder.

DOSSISE VIR RUGSAK SPUIE

PREV-AM®	RUGSAK VOLUME		
Dossise per 100 liters water	10 liter	16 liter	20 liter
100 mℓ	10 mℓ	16 mℓ	20 mℓ
200 mℓ	20 mℓ	32 mℓ	40 mℓ
250 mℓ	25 mℓ	40 mℓ	50 mℓ
500 mℓ	50 mℓ	80 mℓ	100 mℓ

PRODUCT COMPOSITION AND METHOD PATENTED · PRODUKSAMESTELLING EN -METODE GEPATENTEER ZA200405010 / ZA200703565

Oro Agri SA (Pty) Ltd (DIE MAATSKAPPY) WAARBORG DAT DIE PRODUK OOREENSTEM MET SPESIFIKASIES SOOS GESPESEER OP DIE ETIKET EN DAT DIT GEPAS IS VIR DIE BEOOGDE DOEL GENOEM OP DIE ETIKET. DIE MAATSKAPPY VERSKAF GEEN VERDERE DIREKTE OF INDIREKTE WAARBORG NIE DIE MAATSKAPPY BEVEL AN DAT DIE GEBRUIKER DIE NODIGE TOETSE OP 'N GEDEELTE VAN DIE GEWAS DOEN OM DIE GESKIKTHEID VAN DIE PRODUK VIR DIE BEOOGDE GEBRUIK TE BEPAAL.

OUTEURSREG © JULIE 2022 · Oro Agri SA (Pty) Ltd · ALLE REGTE VOORBEHOU · WWW.OROAGRI.CO.ZA · PREV-AM is 'n geregistreerde HANDELSMERK VAN ORO AGRI SA (Pty) Ltd

Witluis*Planococcus ficus*500 ml **PREV-AM**[®]
per 100 l water

0.5% spuitoplossing

- Monitor vir beweging van witluis en begin witluis bespuiting sodra kruipers waargeneem word op lote of blare. Herhaal toediening na 7 dae indien verdere kruiperbeweging waargeneem word
- Miere moet afsonderlik beheer word voordat beheer van witluis gedoen word.
- Let noukeurig op die voorskrifte van die **VERENIGBAARHEID** paragrawe.
- **AANBEVOLE SPUITVOLUME:** 750 l/ha tot 1000 l/ha, afhange van grootte van stokke.

Poeieragtige Skimmel*Uncinula necator*250 - 500 ml **PREV-AM**[®]
per 100 l water

0.25 - 0.5% spuitoplossing

- 'n Program vir die beheer van witroes moet begin sodra die langste lote 'n lengte van 2-5 cm bereik. Gebruik residuele swamdoders tot tydens die blomperiode.
- Die optimale tyd vir gebruik van **PREV-AM**[®] sal wees na blom tot by oes, gedurende die periode wanneer witluis kruiperbeweging verwag sal word (sien witluis aanbeveling).
- Dien **PREV-AM**[®] toe met 7-10 dag intervalle as deel van 'n witroes beheerprogram waarin ander produkte met verskillende metode van werking afgewissel word. **PREV-AM**[®] het 'n nie-spesifieke kontakaksie en kan dus aangewend word in 'n weerstandbestuursprogram teen witroes.
- Gebruik verkieslik 'n siektevoorspellings-model sowel as die siektegeskiedenis van die wingerd as 'n riglyn van die siekerisiko. Indien hierdie inligting 'n lae siektedruk sowel as lae risiko aandui, of geen simptome voorkom nie, gebruik die 0.25% dosis. Wanneer 'n hoër siektedruk en hoër risiko voorkom, of simptome reeds sigbaar is, gebruik die 0.5% dosis.
- Indien toestande gunstig vir infeksie van trosse met groen korrels voorkom, moet **PREV-AM**[®] met 'n geskikte sistemiese swamdoder afgewissel word om die trosse te beskerm.
- **AANBEVOLE SPUITVOLUME:** 500 l/ha tot 1200 l/ha, afhange van grootte van stokke.

Donsskimmel*Plasmopara viticola*500 ml **PREV-AM**[®]
per 100 l water

0.5% spuitoplossing

- 'n Program vir die beheer van Donsskimmel moet begin wanneer die langste lote 'n lengte van 10 cm bereik. Gebruik residuele swamdoders vir voorkomende beheer van donsskimmel.
- Indien **PREV-AM**[®] toegedien word teen 500 l/1000 l vir witluis of witroes, sal hierdie toediening ook donsskimmel beheer wat teenwoordig is tydens die bespuiting.
- Indien gunstige toestande vir infeksie verwag word na toediening van **PREV-AM**[®], moet die produk in mengsel met 'n voorkomende residuele swamdoder toegedien word soos geregistreer.
- Verseker goeie benutting van onderkant van blare.
- **AANBEVOLE SPUITVOLUME:** 500 l/ha tot 1200 l/ha, afhange van grootte van stokke.